

## 2018 S-W Monsoon - Northeast Deficit and Overall Rainfall

### What is the issue?

\n\n

Severe rainfall deficiency in the Northeast has caused an overall deficiency in the country at the end of the current south west monsoon season.

\n\n

### How was rainfall this monsoon?

\n\n

\n

- **Northeast** - Northeast region witnessed a severe rainfall deficiency of 24% this southwest (S-W) monsoon season.

\n

- India Meteorological Department said that it was “very rare” for such a large deficiency in the Northeast.

\n

\n\n

\n

- In the 116-year period from 1901 to 2017, only in 4 years (1992, 2005, 2009 and 2013) has the deficiency in the Northeast exceeded 20%.

\n

\n\n

\n

- Northeast deficiency has caused an overall 9% rainfall deficiency in the country at the end of the season.

\n

\n\n

\n

- **Others Regions** - The distribution of rainfall across Northwest, Central India and Southern Peninsula was “satisfactory”.

\n

- The deficiency in these areas was marginal with 2% each in Northwest and Southern Peninsula, and 7% in Central India.

- \n
- So if the Northeast had received normal rainfall, the all-India rainfall would have been 96% of the Long Period Average.

- \n
- Long Period Average (LPA) is the average annual rainfall for the period 1951-2000.

\n

\n\n

### **What is the cause for deficiency?**

\n\n

- \n
- Starting from the extreme south-western tip of the peninsula, the Southwest monsoon progresses inland.

\n

\n\n

- \n
- But the monsoon trough that usually moves towards the Northeast did not happen this year.
  - Sea surface conditions over the Equatorial Pacific (El Niño conditions) did not have any adverse effect on the rainfall.
  - The IMD has thus planned a detailed scientific analysis on the causes by the end of December 2018.

\n

\n\n

### **Why is 2018 S-W monsoon unique?**

\n\n

- \n
- **Rainfall** - The 2018 monsoon has been characterised by “large day-to-day variability” within the season.
  - Many cities in the country received almost their entire rainfall of the season within a short time.
  - This added up to just a few days to a few weeks within the four-month season (June 1 to September 30).
  - E.g. in 22 cities with sizeable populations, 95% of the monsoon precipitation

occurred over 3 days to 27 days on average.

\n

- Many intense rainfall events during the season have led to flooding like the Kerala floods.

\n

- The season also witnessed formation of a large number of low pressure systems over Bay of Bengal and their westward movement across central India.

\n

- **Water in dams** - The spatial distribution of this year's rainfall is also reflected in the storage capacity of the country's reservoirs.

\n

- Country's major reservoirs at present have 117% of the live storage of the corresponding period last year.

\n

- This is 105% of the average water storage over the last 10 years, as said by the Central Water Commission (CWC).

\n

- Of the five regions, storage in reservoirs in Northern, Eastern, Central and Southern regions is higher than last year.

\n

- Storage in the Western region is less than the storage of last year and also less than the 19-year average storage.

\n

- **Agriculture** - The overall crop acreage during the kharif season is higher by 2.6%.

\n

- This is in comparison to the country's highest ever acreage/record food production during 2017.

\n

- The resultant acreage is largely the result of a good soil moisture distribution across the country.

\n

- Adequate soil moisture available over northern parts of India may help the rabi crop during 2018-19.

\n

\n\n

\n\n

**Source: Indian Express**

\n



**SHANKAR**  
**IAS PARLIAMENT**  
*Information is Empowering*