

## Dual Nutrition Burden in India

### What is the issue?

\n\n

India is witnessing the emerging menace of over-nutrition along with the existing concern of high under-nutrition.

\n\n

### How is under-nutrition in India?

\n\n

\n

- Global Nutrition Report ranks India low in stunting among children aged less than five and in under-5 wasting.

\n

- **Micronutrient deficiency** - The burden of vitamin and mineral deficiencies, called 'hidden hunger', is also considerable.

\n

- This is because a vast majority of Indians eat cereal-based food, mainly wheat and rice.

\n

- There is an insufficient intake of food such as milk, pulses, and fruits and vegetables, which are rich sources of micronutrients.

\n

- **Effects** - Women and children are the most vulnerable to micronutrient deficiencies and consequently have adverse health affects.

\n

- Deficiency of iron in women causes reduced physical work capacity, fatigue, etc.

\n

- Notably it could also lead to depression and post-partum maternal haemorrhage.

\n

- In the case of children, the impact is felt with impairment in growth and cognitive development.

\n

- **Response** - In the last few decades, India has made efforts at improving the food and nutrition security of its population.

- \n
- This was done with strides in technology, irrigation practices, extension services, and progressive agricultural policies.
- \n
- Also, in the 1990s, deficiencies of micronutrients such as zinc, folic acid, magnesium, selenium and vitamin D received more attention.
- \n
- However, despite an overall decrease in cases, the level of under-nutrition still remains high in comparison with world nations.
- \n

\n\n

### **What is the emerging concern?**

\n\n

- \n
- The recent findings of the National Family Health Survey-4 (2015-16) bring out the emerging scenario of over-nutrition in India.
- \n
- With Body Mass Index (BMI) as the measure, the survey identifies-
- \n

\n\n

- \n
- i. around 15% of urban women to be underweight and around 30% of to be overweight or obese
- \n
- ii. around 15% of urban men in the underweight category and a 26% in the overweight and obese category
- \n

\n\n

### **What has led to this?**

\n\n

- \n
- Lifestyle and dietary patterns have undergone dramatic changes, especially among urban sections, in the recent decades.
- \n
- This has contributed to reduction in physical activity and an increasing prevalence of non-communicable diseases.
- \n
- Over-nutrition is traced to be the cause for these diseases due to the imbalance between intake and activity.

\n

- Resultantly, over-nutrition is emerging as a concern among the urban affluent segments.

\n

\n\n

## **What lies ahead?**

\n\n

\n

- India is evidently witnessing a dual nutrition burden.

\n

- The burden of under-nutrition among the poorer sections and that of over-nutrition among the urban well-to-do sections.

\n

- Promotion of appropriate lifestyles and dietary intakes for the prevention and management of over-nutrition and obesity is necessary.

\n

- On the other hand, to ensure food and nutrition security, there is a growing need for a multisectoral approach.

\n

- The policies and programmes of various ministries should be converged for better results in agricultural practices as well as in food styles.

\n

\n\n

\n\n

**Source: The Hindu**

\n\n

\n\n

\n