

Global Nutrition Report 2018

What is the issue?

\n\n

\n

- The Global Nutrition Report 2018 was published recently.
- With India facing a major malnutrition crisis, it calls for concerted actions to address the gaps and concerns.

\n

\n\n

What are the key nutrition indicators?

\n\n

\n

- **Stunting**, or low height for age, is caused by long-term insufficient nutrient-intake and frequent infections.
- **Wasting**, or low weight for height, is a strong predictor of mortality among children under five.
- It is usually the result of acute significant food shortage and/or disease.
- **Overweight** is the condition of having more body fat than is healthy.

\n

\n\n

What are the highlights of the report?

\n\n

\n

- Globally 150.8 million children under five years are stunted and 50.5 million are wasted.
- **Stunting** - Of the three countries that are home to almost half (47.2%) of all stunted children, two are in Asia.

\n

- Of these, with 46.6 million (31%) children who are stunted, India tops the list of countries.
\n
- It is followed by Nigeria (13.9 million) and Pakistan (10.7 million).
\n
- **Wasting** - More than half of the world's children impacted by wasting (26.9 million) live in South Asia.
\n
- India accounted for 25.5 million children who are wasted, followed by Nigeria (3.4 million) and Indonesia (3.3 million).
\n
- **Overweight** - India also figures among the set of countries that has more than a million overweight children.
\n
- The other nations are China, Indonesia, Egypt, US, Brazil and Pakistan.
\n
- In four countries, more than a fifth of all children are overweight - Ukraine, Albania, Libya and Montenegro.
\n
- Of the around 38 million children globally overweight, some 5.4 million are in South Asia and 4.8 million are in East Asia.
\n
- Prevalence of overweight children is the highest in upper-middle income countries and the lowest in low-income countries.
\n
- In urban areas, there are 7.1% overweight children on average, whereas in rural areas 6.2% children are overweight.
\n
- Overweight is slightly more common among boys (6.9%) than girls (6.1%).
\n
- As with obesity, among adults, women are more overweight than among men (39.2% and 38.5% respectively in 2016).
\n
- Conversely, diabetes is more common among men than women (9% and 7.9% respectively in 2014).
\n
- The health consequences of being overweight and obese contribute to an estimated four million deaths globally.
\n
- **Malnutrition** - Malnutrition is responsible for more ill-health than any other cause.
\n
- Of the 141 countries analysed, 88% of countries experience more than one form of malnutrition.

- \n
- The problem of malnutrition remains severe across all regions but none of the countries are on course to meet all nine global nutrition targets.
- \n
- **Packaged foods** - As per the report, only 21% of packaged foods in India are rated as being healthy.
- \n
- This is based on overall energy, salt, sugar and saturated fat on the negative side, and vegetable, fruit, protein, fibre and calcium as positive factors.
- \n

\n\n

What is the case with malnutrition in India?

\n\n

- \n
- **Spatial variation** - An International Food Policy Research Institute (IFPRI) study found that stunting varied greatly from district to district (nearly 12% to 65%).
- \n
- Notably, 239 of 604 districts have stunting levels above 40%.
- \n
- There are high and very high levels of stunting mainly in central and northern India (more than 30% and 40%, respectively).
- \n
- In contrast, less than 20% is the condition in almost the entire south.
- \n
- **Factors** - Factors associated with gender, education, economic status, health, hygiene, and other demographic factors explain the spatial variation.
- \n
- Factors such as women's low BMI accounted for 19% of the difference between the low versus high-burden districts.
- \n
- Other influential gender-related factors included maternal education (accounted for 12%), age at time of marriage (7%) and antenatal care (6%).
- \n
- Children's diets (9%), assets (7%), open defecation (7%) and household size (5%) were also influential.
- \n

\n\n

What should be done?

\n\n

\n

- The district- and region-wise sharp contrast shows the important role played by political commitment and administrative efficiency.

\n

- It also reflects the role of literacy and women's empowerment in ensuring children's health.

\n

- The figures thus call for immediate policy action to reduce inequalities and childhood stunting.

\n

\n\n

\n

- With district-wise data, state governments should address the determining factors for malnutrition.

\n

- Food and freedom go together, and the availability of one strongly influences access to the other.

\n

- So, social institutions can work to improve nutrition and children's welfare in free societies.

\n

- Governments should acknowledge the linkages and commit themselves to improved nutritional policies.

\n

- In states where the Anganwadi Services scheme does not work well, it must be subjected to a rigorous review, and targeted interventions for supplementary nutrition should be made.

\n

\n\n

\n\n

Source: Economic Times, The Hindu

\n

