

## **LIGO to Publish Paper on Analysis Techniques**

### **Why in news?**

\n\n

\n

- The LIGO (Laser Interferometer Gravitational-Wave Observatory) Scientific Collaboration detected gravitational waves in 2015.

\n

- It has recently announced that it would publish a detailed explanation of how it analyses the noise in its detectors.

\n

\n\n

### **What was the 2015 discovery?**

\n\n

\n

- Gravitational waves are ripples in the fabric of space-time, arising from the merger of a pair of black holes in distant space.

\n

- LIGO's 2015 announcement of the discovery of gravitational waves was an exciting finding in physics for decades.

\n

- The discovery confirmed a prediction made by Einstein.

\n

- It stated that space-time itself can squeeze and stretch in rhythmic waves, when deformed by cataclysmic events like collision of black holes.

\n

- The collaboration's founders were awarded the Nobel prize in physics in 2017. Click [here](#) to know more.

\n



\n\n

## WHAT IS LIGO?

The advanced Laser Interferometer Gravitational Wave Observatory (or LIGO) is at the centre of the path-breaking find:

**The LIGO experiment is an example of extreme engineering chasing an impossible dream**

The twin LIGO installations – one in Livingston, Louisiana, and the other in Hanford, Washington – are located 3,000km apart



- ▶ Built **3,000km** apart, operating in unison
- ▶ To make the smallest measurement ever attempted by science – **a motion 10,000 times smaller than an atomic nucleus**
- ▶ Caused by the most violent and cataclysmic events in the Universe occurring millions of light years away
- ▶ Can detect gravitational waves in a volume of 1 billion cubic light years – **covering about 1 million Milky Way type galaxies**

- ▶ To detect a gravitational wave we should be able to tell when something changes in length by a few parts in 10 to the power 23
- ▶ LIGO makes the smallest measurement ever attempted – a motion 10,000 times smaller than an atomic nucleus
- ▶ It's like trying to hear a song being hummed in a very, very noisy party

\n\n

## What were the further observations?

\n\n

\n

- Since detecting the binary black hole (BBH) merger, the LIGO Scientific Collaboration (LSC) has made six such observations.

\n

- Five of these were mergers of black holes in very different locations in space and with very different characteristics such as mass.

\n

- Another was the merger of a pair of so-called neutron stars (binary neutron stars).


\n

- The last few detections have been done in conjunction with another detector,

Virgo (Italy-based).

\n

\n\n



\n\n

### What is the need for LIGO's explanation now?

\n\n

\n

- **Challenge** - LIGO's detectors aim to measure a shortening of space equivalent to about a thousandth of the width of a proton.
- This sort of measurement is swamped by natural thermal vibrations, known as noise.
- This makes picking out the signal from a gravitational wave tricky and challenging.
- The collaboration thus used sophisticated analysis techniques to remove this noise.
- Also, after the first discovery, the LSC made public its data on these techniques.

\n

\n\n

\n

- **Dispute** - Analysing the data, in 2017, a group of scientists questioned the validity of the first detection.
- Weeding out noise from the signal is crucial in any such experiment.

\n

\n

- Some claimed that this had not been done properly by the LSC.

\n

- They argued that the two detectors belonging to LIGO were correlated and that this led to a correlation in the noise factor.

\n

\n\n

\n

- Other scientific investigations also uncovered a number of irregularities in the data.

\n

- **LSC** - After a long silence, recently, the LSC has thus put up a clarification on its website.

\n

- The LIGO collaboration is learnt to be in the process of preparing their paper clarifying their approach and explaining the analysis techniques.

\n

\n\n

\n\n

**Source: The Hindu, NewScientist**

\n

