

## Neglected Tropical Diseases in South Asian Region

### What is the issue?

\n\n

\n

- South Asian region has recorded a notable progress in eliminating and containing various diseases.

\n

- However, the status of neglected tropical diseases (NTDs) in the South Asian region calls for a last-mile push.

\n

\n\n

### What are the health achievements in South Asia?

\n\n

\n

- Maternal and neonatal tetanus has been eliminated.

\n

- HIV, TB and malaria epidemics have been halted and reversed after decades of struggle.

\n

- These will hopefully be ended altogether in the coming years.

\n

- **Regional** - India is now yaws-free.

\n

- Maldives, Sri Lanka and Thailand have eliminated *lymphatic filariasis* as a public health problem.

\n

- Nepal is in the process of validating the elimination of *trachoma* as a public health problem.

\n

- Nepal also maintained the elimination target for *kala-azar* for more than 3 consecutive years.

\n

- By the end of 2017, 100% of sub-districts in Bangladesh and 90% of blocks in India had done the same.

\n

- As a whole, the region remains responsible for the world's largest *preventive*

*chemotherapy campaign.*

\n

\n\n

## **What are the shortfalls?**

\n\n

\n

- NTDs include diseases like leprosy, trachoma, lymphatic filariasis and kala-azar.

\n

- As the name suggests, NTDs take their most severe toll on the poor and marginalised communities.

\n

- The political influence of these sections is limited and their health needs are often overlooked.

\n

- Evidently, South Asian region has eliminated leprosy as a public health problem.

\n

- But the disease continues to circulate among vulnerable communities.

\n

- They accounts for more than 60% of leprosy-caused grade two disabilities worldwide.

\n

- Similarly, several member countries have eliminated *lymphatic filariasis*.

\n

- However, its burden continues to haunt communities in remote and hard-to-reach areas elsewhere.

\n

- Region-wide, 53% of all people require mass drug administration to stay free of this disfiguring disease.

\n

\n\n

## **What is the way forward?**

\n\n

\n

- No member country of the South Asian region can be complacent as one of the 20 NTDs is endemic to each of them.

\n

- Stable **domestic funding** for national programmes is central to taking forward the progress achieved so far.

\n

- Member countries must ensure programming is both dynamic and flexible i.e. pursuing large-scale campaigns as well as working creatively across sectors.

\n

- Going for **mass drug administration** where appropriate and strengthening NTD-related services at the primary level are essential.

\n

- Simultaneously, **specific efforts** such as roping in agro-engineers in the battle against *schistosomiasis* can be taken up.

\n

- Similarly, **vector control** must be actively pursued and harnessed by programmes at the local level.

\n

- These can have immediate and substantial impact on disease transmission in NTD-affected areas.

\n

- Importantly, member countries should take full advantage of innovations in **research and technology**.

\n

- The effective use of rapid **diagnostics** can facilitate swift and accurate diagnosis of a range of NTDs.

\n

- IT infrastructure should be integrated with existing surveillance systems to allow programme managers to gather, analyse and act on real-time data.

\n

- Ensuring all communities have **access** to health services and the benefits they provide is essential to make them free of NTDs.

\n

\n\n

\n\n

**Source: Business Line**

\n\n

**Quick Fact**

\n\n

**Preventive Chemotherapy Campaign**

\n\n

\n

- This is a public health strategy recommended by WHO against a set of Neglected Tropical Diseases.

\n

- Preventive Chemotherapy consists of the regular, large-scale administration of drugs, either alone or in combination.

\n

- It is administered to entire population groups, with the aim of reducing transmission and associated morbidity.

\n

\n\n

## **Lymphatic filariasis**

\n\n

\n

- Lymphatic filariasis is commonly known as elephantiasis.

\n

- It is caused by infection with parasites classified as nematodes (roundworms), which are transmitted to humans through mosquitoes.

\n

- Mosquitoes are infected with microfilariae by ingesting blood when biting an infected host.

\n

- Infection is usually acquired in childhood causing hidden damage to the lymphatic system.

\n

- Its visible manifestations may occur later in life, causing temporary or permanent disability.

\n

\n\n

## **Schistosomiasis**

\n\n

\n

- Schistosomiasis is an acute and chronic disease caused by parasitic worms.

\n

- People are infected during routine agricultural, domestic, occupational, and recreational activities, which expose them to infested water.

\n

- Water contamination due to excreta containing parasite eggs, which hatch in water, is also a mode of transmission.

\n

\n\n

## **Trachoma**

\n\n

- \n
  - Trachoma is a disease of the eye caused by infection with the bacterium *Chlamydia trachomatis*.
- \n
  - It causes blindness or visual impairment. It is the cause for about 1.4% of all blindness worldwide.
- \n
  - Blindness from trachoma is irreversible.
- \n
  - Environmental risk factors influencing the transmission of the disease include:
- \n

\n\n

- \n
  - i. poor hygiene
- \n
  - ii. crowded households
- \n
  - iii. water shortage
- \n
  - iv. inadequate latrines and sanitation facilities
- \n

\n\n

## **Kala-azar**

\n\n

- \n
  - Visceral leishmaniasis (VL), also known as kala-azar, is caused by the protozoan *Leishmania* parasites.
- \n
  - It is transmitted to humans through infected sandflies.
- \n
  - The parasite migrates to the internal organs such as the liver, spleen, and bone marrow; if left untreated, it may result in death.
- \n
  - It is characterized by irregular bouts of fever, weight loss, enlargement of the spleen and liver, and anaemia.
- \n

- It is endemic to the Indian subcontinent in 119 districts in four countries (Bangladesh, Bhutan, India and Nepal).

\n

\n

