

## **Northeast Cancer challenge**

### **What is the issue?**

\n\n

\n

- Lifestyle-related cancers top the chart in the northeast region.

\n

- This has an overall high incidence of cancer in India.

\n

\n\n

### **What is the status of cancer in north east?**

\n\n

\n

- A recent report was compiled by the National Centre for Disease Informatics and Research under the Indian Council of Medical Research (ICMR).

\n

- Nearly 4% of the cancer cases detected in Mumbai's Tata Memorial Hospital (TMH) are in patients from the northeast.

\n

- It shows Oesophageal cancer is the highest in men followed by cancers of the lung and stomach.

\n

- It also analyses a total of 37,448 cancer cases reported from 11 population-based cancer registries in the States of the northeast from 2012 to 2014.

\n

- Together, they form 33.5% of all cancers in the northeast among men.

\n

- In women, breast cancer is the highest followed by cervix and oesophageal cancer (33.8%).

\n

\n\n

### **What are the reasons of the complications?**

\n\n

\n

- A feature that stands out in Arunachal Pradesh, Assam, Manipur, Meghalaya, Mizoram, Nagaland, Sikkim and Tripura is excessive tobacco consumption.  
\n
- People in the region are also very addicted to raw betel nut and its fermented variety.  
\n
- Both tobacco and betel nuts are known carcinogenic substances.  
\n
- **Oesophageal cancer** is also linked to consuming extremely hot beverages.  
\n
- People who gulped down very hot tea (simmering temperature) were four times more prone to oesophageal cancer.  
\n
- There are certain dietary habits that definitely play a role in the increased incidence of cancer in the region.  
\n
- The **genetic variations** as a result of there being a large number of ethnic groups are also a contributing factor.  
\n
- **Stomach cancer** is linked to H. pylori bacteria infection that happens due to contaminated food and water, obesity, and a diet consisting of salty or smoked foods.  
\n
- **Breast cancer** too is linked with obesity and late pregnancies among other risk factors.  
\n
- In **cervical cancer**, the most important risk factor is a Human Papilloma Virus (HPV) infection that is mainly transmitted through sexual intercourse.  
\n
- Multiple sex partners (polyandries and polygamies societies are prevalent in North east region), a lack of hygiene and smoking are some factors linked to the spread of cervical cancers.  
\n

\n\n

### **What are the challenges patients are facing?**

\n\n

- Most cancer patients have to travel outside the northeast for treatment and care due to an inadequate number of cancer facilities.  
\n
- There are only about eight centres that offer various types of cancer care such as surgery, chemotherapy and radiation therapy and most of them are

in Assam.

\n

- Cancer cannot be treated from afar as therapy is a prolonged one.
- Travelling in the region is not easy due to the difficult terrain and this results in a low compliance of treatment.

\n

\n\n

\n\n

**Source: The Hindu**

\n

