

## **Patient Suicides - The way ahead**

### **What is the issue?**

\n\n

\n

- A senior IPS officer had recently committed suicide, due to depression caused by a prolonged chronic illness.

\n

- Monitoring the mental health of patients is vital to avoid such episodes.

\n

\n\n

### **What is the state of patients committing suicide?**

\n\n

\n

- Family problems account for the most number of suicides in India, accounting for 27.6% of all suicides.

\n

- This is followed by illness driven suicides which stands at a whopping 15.8%.

\n


- Hence, sad moods or odd behaviour should not be brushed aside as a “normal reaction” to being ill and should be seen as possible symptoms of depression.

\n

- Doctors and family need to keep a close eye on symptoms of depression in the patients who happen suffer chronic or long lasting disorders/diseases.

\n

\n\n



\n\n

## What drives suicides?

\n\n

\n

- Most patients who commit suicide aren't at their terminal stage but the ones who can't mentally cope with the initial stages of the illness.

\n

- Patients suffering from a prolonged illness are affected as they miss out on a lot in life, which is taken for granted by others.

\n

- They have to make many lifestyle changes and simultaneously cope with the side effects of medication.

\n

- These are highly stressful for a vast majority of the patients as they find it difficult to make peace with their new situation.

\n

- In addition to this, some also have to go through eternal chronic pain, which is the main reason that drives them to suicide.

\n

\n\n

## How do we address this problem?

\n\n

\n

- While illness is what causes depression usually, in considerable number of cases, it is the medication that creates depression.

\n

- Continuous counselling and keeping a close eye on depressive symptoms in patients is the key to preventing illness driven suicides.

\n

- Counselling should not be isolated as a psychiatric measure, but should rather be incorporated to an extent within the conventional treatment

centres.

\n

- All physicians and medical staff treating such patients need to be able to pick up depressive symptoms through a patient's mannerism.

\n

\n\n

\n\n

**Source: The Hindu**

\n

