

## Three new additions to the vaccination basket

### Why in news?

\n\n

\n

- The basket of vaccines in India's Universal Immunisation Programme (UIP) was static for many years — until the entry of the Pentavalent vaccine [Diphtheria, Tetanus, Pertussis, Hepatitis B, and Haemophilus B (HIB)], and Japanese Encephalitis vaccines.

\n

- There have been regular additions to the basket since. Two new vaccines — **Measles-Rubella (MR) and Pneumococcal Conjugate** — are lined up for launch in January and February respectively, and a **third, Rotavirus Vaccine**, will become part of the UIP in 5 states — from February.

\n

\n\n

### What is rubella, and why is vaccination for it important?

\n\n

\n

- More commonly known as German Measles, **Congenital Rubella Syndrome, or CRS**, is believed to affect about 25,000 children born in India every year.

\n

- Symptoms can include cataracts and deafness, and the disease can also affect the heart and the brain.

\n

- 10-30% of adolescent females and 12-30% of women in the reproductive age-group are susceptible to rubella infection in India.

\n

\n\n

### In which states will the MR vaccine be introduced?

\n\n

\n

- After its formal launch on January 17, the vaccine will be introduced in **Goa**,

**Karnataka, Lakshadweep, Puducherry and Tamil Nadu** around February 3-4.

\n

- The Health Ministry will run a campaign among children aged 9 months to 15 years before making the vaccine a part of routine immunisation. Two shots will be given — one between the ages of 9-12 months, the other at age one-and-a-half.

\n

- The monovalent measles vaccine is already part of the UIP basket of 10 vaccines; it will be discontinued once MR is introduced.

\n

\n\n

### **What is Pneumococcal Conjugate Vaccine (PCV)?**

\n\n

\n

- PCV is a **mixture of several bacteria of the pneumococci family**, which are known to cause pneumonia — hence ‘conjugate’ in the name. Pneumonia caused by the pneumococcus bacteria is supposed to be the most common.

\n

- Pneumonia and diarrhoea have long been responsible for the most child deaths in India — approximate estimates say **pneumonia is responsible for about 20% of under-5 child mortality in India**, of which half are of pneumococcal origin.

\n

- In 2008, the WHO’s Child Health Epidemiology Reference Group reported that 5 countries in which 44% of the world’s children aged less than 5 years live (India, China, Pakistan, Bangladesh, Indonesia and Nigeria) contribute more than half of all new pneumonia cases annually.

\n

- It estimated around 43 million pneumonia cases (23% of the global total) and an incidence of 0.37 episodes per child-year for clinical pneumonia in India.

\n

\n\n

### **Which states will PCV be introduced?**

\n\n

\n

- In Himachal Pradesh and parts of Uttar Pradesh and Bihar from March 17. Three doses will be administered at one-and-a-half months, three-and-a-half months and 9 months.

- \n
- The annual incidence of severe pneumococcal pneumonia in India was estimated to be **4.8 episodes per 1,000 children younger than 5 years.**
  - The top five contributors to India's pneumococcal pneumonia burden in terms of number of cases and deaths were Uttar Pradesh, Bihar, Madhya Pradesh, Rajasthan and Jharkhand.
- \n

\n\n

### **Where will the rotavirus vaccine be introduced?**

\n\n

- \n
- **Rotavirus infections are the most common cause of diarrhoea in children.** The rotavirus vaccine first became a part of UIP in April 2016.
  - An estimated 1 lakh children die every year of the disease.
  - The vaccine is currently being administered in HP, Haryana, Odisha and AP. From February, it will be a part of UIP also in Assam, Tripura, Rajasthan, Madhya Pradesh and Tamil Nadu.
- \n

\n\n

\n\n

**Category: Prelims & Mains | GS - II | Health**

\n\n

**Source: The Indian Express**

\n

